

Information for the User

EROPID
SILDENAFIL

Eropid 50 mg film-coated tablets (Sildenafil Citrate)



1. What Eropid is and what it is used for

Eropid belongs to a group of medicines called phosphodiesterase type 5 (PDE5) inhibitors. It works by helping to relax the blood vessels in your penis, allowing blood to flow into your penis when you get sexually excited. Eropid will only help you to get an erection if you are sexually stimulated. You should not take Eropid if you do not have erectile dysfunction. You should not take Eropid if you are a woman.

Eropid is a treatment for men with erectile dysfunction, sometimes known as impotence. This is when a man cannot get, or keep a hard, erect penis suitable for sexual activity.

2. What you need to know before you take Eropid

Do not take Eropid

- If you are allergic to Sildenafil or any of the other ingredients of this medicine (listed in section 6).
- If you are taking medicines called nitrates, as the combination may cause a potentially dangerous decrease in your blood pressure. Tell your doctor if you are taking any of these medicines, which are often given for relief of angina pectoris (or "chest pain"). If you are not certain, ask your doctor or pharmacist.
- If you are using any of the drugs known as nitric oxide donors such as amyl nitrite ("poppers"), as the combination may also lead to a potentially dangerous decrease in your blood pressure.
- If you are allergic (hypersensitive) to Sildenafil or any of the other ingredients of Sildenafil.
- If you have a severe heart or liver problem.

- If you have recently had a stroke or a heart attack, or if you have low blood pressure.
- If you have certain rare inherited eye diseases (such as retinitis pigmentosa).
- If you have ever had loss of vision due to non-arteritic anterior ischaemic optic neuropathy (NAION).

Warnings and precautions

Talk to your doctor before taking Eropid

- If you have sickle cell anaemia (an abnormality of red blood cells), leukaemia (cancer of blood cells), multiple myeloma (cancer of bone marrow).
- If you have a deformity of your penis or Peyronie's Disease.
- If you have problems with your heart. Your doctor should in that case carefully check whether your heart can take the additional strain of having sex.
- If you currently have a stomach ulcer, or a bleeding problem (such as haemophilia).
- If you experience sudden decrease or loss of vision, stop taking Eropid and contact your doctor immediately.

You should not use Eropid with any other oral or local treatment for erectile dysfunction.

Children and adolescents

Eropid should not be given to individuals under the age of 18.

Patients with kidney or liver problems

You should tell your doctor if you have kidney or liver problems. Your doctor may decide on a lower dose for you.

Other medicines and Eropid

Tell your doctor or pharmacist if you are taking , have recently taken or might take any other medicines.

Eropid tablets may interfere with some medicines, especially those used to treat chest pain. In the event of a medical emergency, you should tell any health care professional treating your condition that you have taken Eropid and when you did . Do not take Eropid with other medicines unless your doctor tells you that you can.

You should not take Eropid if you are taking medicines called nitrates, as the combination of these products may cause a potentially dangerous decrease in your blood pressure. Always tell your doctor or pharmacist if you are taking any of these medicines that are often used for the relief of angina pectoris (or "chest pain").

You should not take Eropid if you are using any of the drugs known as nitric oxide donors such as amyl nitrite ("poppers"), as the combination may lead to a potentially dangerous decrease in your blood pressure.

If you are taking medicines known as protease inhibitors, such as for the treatment of HIV, your doctor may start you on the lowest dose (25 mg) of Eropid.

Some patients who take alpha-blocker therapy for the treatment of high blood pressure or prostate enlargement may experience dizziness or light-headedness, which may be caused by low blood pressure upon sitting or standing up quickly. Certain patients have experienced these symptoms when taking Eropid with alpha-blockers . This is most likely to occur within 4 hours after taking Eropid. In order to reduce the likelihood that these symptoms occur, you should be on a regular daily dose of your alpha-blocker before you start Eropid. Your doctor may start you on a lower dose (25 mg) of Eropid.

Eropid with food, drink and alcohol

Eropid can be taken with or without food. However, you may find that Eropid takes longer to start working if you take it with a heavy meal.

Drinking alcohol can temporarily Impair your ability to get an erection. To get the maximum benefit from your medicine, you are advised not to drink excessive amounts of alcohol before taking Eropid.

Pregnancy and breast-feeding

Eropid is not indicated for use by women.

Driving and using machines

Eropid can cause dizziness and can affect vision. You should be aware of how you react to Sildenafil before you drive or use machinery.

Eropid contains Tartrazine

May cause allergic reactions as it contains Tartrazine (E-102). It can worsen asthma.

3. How to take Eropid

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The usual starting dose is 50 mg.

You should not take Eropid more than once a day.

You should take Eropid about one hour before you plan to have sex. Swallow the tablet whole with a glass of water.

If you have the Impression that the effect of Eropid is too strong or too weak, talk to your doctor or pharmacist.

Eropid will only help you to get an erection if you are sexually stimulated. The amount of time Eropid takes to work varies from person to person, but it normally takes between half an hour and one hour. You may find that Eropid takes longer to work If you take It with a heavy meal.

If Eropid does not help you to get an erection, or if your erection does not last long enough for you to complete sexual Intercourse you should tell your doctor.

If you take more Eropid than you should:

You may experience an increase in side effects and their severity. Doses above 100 mg do not increase the efficacy.

You should not take more tablets than your doctor tells you to.

Contact your doctor if you take more tablets than you should.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects although not everybody gets them. The side effects reported in association with the use of Eropid are usually mild to moderate and of short duration.

If you have chest pains during or after intercourse:

- Get in a semi-sitting position and try to relax.
- Do not use nitrates to treat your chest pain.
- Contact your doctor immediately.

All medicines including Eropid can cause allergic reactions. You should contact your doctor immediately if you experience any of the following symptoms after taking Eropid: sudden wheeziness, difficulty in breathing or dizziness, swelling of the eyelids, face, lips or throat.

Prolonged and sometimes painful erections have been reported after taking

Eropid. If you have an erection which lasts for more than 4 hours, you should contact a doctor immediately.

If you experience a sudden decrease or loss of vision, stop taking Eropid and contact your doctor immediately.

A very common side effect (may affect more than 1 in 10 people) is headache.

Common side effects (may affect up to 1 in 10 people) include: facial flushing, indigestion, effects on vision including colour tinge to vision, light sensitivity, blurred vision or reduced sharpness of vision) stuffy nose and dizziness.

Uncommon side effects (may affect up to 1 in 100 people) include: vomiting, skin rash, bleeding at the back of the eye, eye irritation, bloodshot eyes/red eyes, eye pain, double vision, abnormal sensation in the eye, watery eyes, irregular or rapid heartbeat, muscle pain, feeling sleepy, reduced sense of touch, vertigo, ringing in the ears, nausea, dry mouth, penile bleeding, presence of blood in semen and/or urine, chest pain and feeling tired.

Rare side effects (may affect up to 1 in 1,000 people) include: high blood pressure, low blood pressure, fainting, stroke, heart attack, nosebleed, sudden decrease or loss of hearing and non-arteritic anterior ischaemic optic neuropathy.

Additional side effects reported from post-marketing experience include: pounding heartbeat, chest pain, sudden death, or temporary decreased blood flow to parts of the brain. Most, but not all, of these men had heart problems before taking this medicine. It is not possible to determine whether these events were directly related to Eropid. Cases of convulsions or seizures and serious skin reactions characterised by rash, blister, peeling skin and pain which require immediate medical attention have also been reported.

Reporting of a side effect

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme. Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google

Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.



5. How to store Eropid

Keep this medicine out of the sight and reach of children.

This medicinal product does not require any special temperature storage conditions.

Do not use this medicine after the expiry date which is stated on the carton.

The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Eropid contains

The active substance is Sildenafil.

Each film-coated tablet contains 50 mg of Sildenafil, as the citrate.

The other ingredients are:

- Tablet core; cellulose powder, colloidal silicon (anhydrous), calcium phosphate dibasic (anhydrous), croscarmellose sodium, magnesium stearate,
- Film coat: hypromellose, titanium dioxide (E171), triacetin, indigotine, aluminium lake (E132), brilliant blue aluminium lake (E133), tartrazine, aluminium lake (E102)

What Eropid looks like and contents of the pack

Sildenafil film-coated tablets are blue, circular and biconvex tablets. The tablets are provided in blister packs containing 2, 4, 8 or 12 tablets. Not all pack sizes may be marketed